

**M -theory on $AdS_4 \times Q^{11}$:
the complete $Osp(2|4) \times SU(2) \times SU(2) \times SU(2)$ spectrum
from harmonic analysis [†]**

Paolo Merlatti

*Dipartimento di Fisica Teorica, Università di Torino
and I.N.F.N., sezione di Torino
via P. Giuria 1, I-10125 Torino, Italy
Paolo.Merlatti@to.infn.it*

Abstract

In this paper by means of harmonic analysis we derive the complete spectrum of $Osp(2|4) \times SU(2) \times SU(2) \times SU(2)$ multiplets that one obtains compactifying $D = 11$ supergravity on the homogeneous space Q^{11} .

In particular we analyze the structure of the short multiplets and compare them with the corresponding composite operators of the $\mathcal{N} = 2$ conformal field theory dual to such a compactification, found in a previous publication. We get complete agreement between the quantum numbers of the supergravity multiplets on one side and those of the conformal operators on the other side, confirming the structure of the conjectured *SCFT*.

However the determination of the actual spectrum by harmonic analysis teaches us a lot more: indeed we find out which multiplets are present for each representation of the isometry group, how many they are, the exact values of the hypercharge and of the “energy” for each multiplet.

[†] Research partially supported by the EC RTN programme HPRN-CT-2000-00131

1 Introduction

The basic principle of the *AdS/CFT* correspondence [2, 3, 4] states that every consistent *M*-theory or type II background with metric $AdS_{p+2} \times M^{d-p-2}$ in d -dimensions, where M^{d-p-2} is an Einstein manifold, is associated with a conformal quantum field theory living on the boundary of AdS_{p+2} . The background is typically generated by the near horizon geometry of a set of p -branes and the boundary conformal field theory is identified with the IR limit of the gauge theory living on the world-volume of the p -branes.

In this paper we focus on the case $p = 2$ when M is a coset manifold G/H with $\mathcal{N} = 2$ supersymmetry. In this case one remarkable example with a non-trivial smooth manifold $M^7 = M^{111}$ was found in [1, 5]. Some general properties and the complete spectrum of the M^{111} compactification have been discussed in [5], while in [1] the associated superconformal theory that matches supergravity predictions has been identified.

For the manifold $M^7 = Q^{111}$ an ansatz for the superconformal field theory was given in [1]¹. In this paper we study $D = 11$ supergravity compactified on Q^{111} . So in this case we reverse the usual procedure and we rather use the SCFT side to make prediction about the Kaluza-Klein spectrum. We show that all the predictions are completely verified.

Actually this paper constitutes a step of a wider project which deals with the specific case of $D = 11$ supergravity compactified on internal homogeneous seven-manifolds: $M^7 = (G/H)^7$. Many examples were studied in the old days of Kaluza Klein theories [6] (see [7, 9, 10, 11, 12, 13, 14, 15, 16] for the cases S^7 and squashed S^7 , see [17, 18, 19, 20, 21, 23] for the case $M^{p,q,r}$, see [24, 25] for the case Q^{111} , see [16, 19, 26, 27, 28] for general methods of harmonic analysis in compactification and the structure of $Osp(N|4)$ supermultiplets, for an updated recent review see [22], and finally see [29] for a complete classification of G/H compactifications.)

This paper is organized as follows. In section two we briefly review the essential features of Q^{111} geometry. In section three, using standard techniques of harmonic analysis, we compute the spectrum of the seven-dimensional Laplace-Beltrami operators that we need to calculate the complete mass-spectrum of the four-dimensional fields. Finally in section four we organize all the fields in multiplets of the relevant superalgebra.

2 The differential geometry of Q^{111}

Let us briefly review the essential features of Q^{111} geometry. For notations and basic lore we mainly refer to [18, 19, 30].

The homogeneous space Q^{111} is the quotient of $G = SU(2) \times SU(2)' \times SU(2)'' \times U(1)$ by the action of its subgroup $H = U(1)' \times U(1)'' \times U(1)'''$, where the embedding of H in G is defined as follows. The H subgroup is generated by $Z^{(1)}$, $Z^{(2)}$ and $Z^{(3)}$, being three independent linear combinations of the four abelian generators of G . Using the Pauli matrices for $SU(2)$ and being K the generator of the $U(1) \in G$ these four generators are $J_3 = -\frac{i}{2}\sigma_3$, $J_{3'} = -\frac{i}{2}\sigma'_3$, $J_{3''} = -\frac{i}{2}\sigma''_3$ and K . The three linear combinations $Z^{(1)}$, $Z^{(2)}$ and $Z^{(3)}$ are defined as a

¹A first attempt at the derivation of such a superconformal field theory dual of Q^{111} was put forward in [8]

basis for the orthogonal complement of the generator

$$Z = -\frac{1}{\sqrt{3}} \left(\frac{i}{2}\sigma_3 + \frac{i}{2}\sigma'_3 + \frac{i}{2}\sigma''_3 \right) \quad (2.1)$$

A basis for the three abelian generators of H is then given by

$$Z^{(1)} = \frac{1}{\sqrt{3}} \left(-\frac{i}{2}\sigma_3 + \frac{i}{2}\sigma'_3 \right) \quad (2.2)$$

$$Z^{(2)} = \frac{1}{\sqrt{3}} \left(-\frac{i}{2}\sigma_3 - \frac{i}{2}\sigma'_3 + 2\frac{i}{2}\sigma''_3 \right) \quad (2.3)$$

$$Z^{(3)} = -\frac{1}{\sqrt{3}} \left(\frac{i}{2}\sigma_3 + \frac{i}{2}\sigma'_3 + \frac{i}{2}\sigma''_3 + 3K \right) \quad (2.4)$$

From an explicit parametrization of the coset G/H we can construct the left-invariant one-forms on G/H as:

$$\Omega(y) = L^{-1}(y)dL(y) = \Omega^\Lambda(y)T_\Lambda, \quad (2.5)$$

which satisfy the Maurer-Cartan equations

$$d\Omega^\Lambda + \frac{1}{2}\mathcal{C}^\Lambda_{\Sigma\Pi}\Omega^\Sigma \wedge \Omega^\Pi = 0 \quad (2.6)$$

with the structure constants of G :

$$[T_\Sigma, T_\Pi] = \mathcal{C}^\Lambda_{\Sigma\Pi}T_\Lambda. \quad (2.7)$$

The one-forms Ω^Λ can be separated into a set $\{\Omega^H\}$ corresponding to the generators of the subalgebra $\mathbb{H}$ and a set $\{\Omega^\alpha\}$ corresponding to the coset generators. These latter can be identified with the $SU(2) \times SU(2)' \times SU(2)''$ invariant seven-vielbeins on G/H :

$$\begin{aligned} \mathcal{B}^\alpha &\equiv (\mathcal{B}^i, \mathcal{B}^{i'}, \mathcal{B}^{i''}, \mathcal{B}^z), \\ \left\{ \begin{array}{l} \mathcal{B}^i &= \frac{\sqrt{2}}{8}\Omega^i, \\ \mathcal{B}^{i'} &= \frac{\sqrt{2}}{8}\Omega^{i'}, \\ \mathcal{B}^{i''} &= \frac{\sqrt{2}}{8}\Omega^{i''} \\ \mathcal{B}^z &= \frac{1}{8}(\Omega^0 + \Omega^{0'} + \Omega^{0''}) = \frac{\sqrt{3}}{8}\Omega^z, \end{array} \right. \quad (2.8) \end{aligned}$$

where the multiplicative coefficients in front of the vielbeins have been properly chosen to let the metric on G/H be Einstein.

The invariant forms Ω^H are:

$$\left\{ \begin{array}{l} \Omega^{Z^{(1)}} = \frac{\sqrt{3}}{2}(\Omega^0 - \Omega^{0'}), \\ \Omega^{Z^{(2)}} = \frac{\sqrt{3}}{6}(\Omega^0 + \Omega^{0'} - 2\Omega^{0''}). \end{array} \right. \quad (2.9)$$

The spin-connection $\mathcal{B}^\alpha_\beta$ is easily determined from the vielbeins $\mathcal{B}^\alpha$ by imposing vanishing torsion:

$$d\mathcal{B}^\alpha - \mathcal{B}^\alpha_\beta \wedge \mathcal{B}^\beta = 0, \quad (2.10)$$

$$\begin{cases} \mathcal{B}^{ij} &= \frac{1}{\sqrt{3}}\epsilon^{ij}\left(\frac{2\sqrt{3}}{3}\mathcal{B}^z + \mathcal{B}^{z(1)} + \mathcal{B}^{z(2)}\right), \\ \mathcal{B}^{i'j'} &= \frac{1}{\sqrt{3}}\epsilon^{i'j'}\left(\frac{2\sqrt{3}}{3}\mathcal{B}^z - \mathcal{B}^{z(1)} + \mathcal{B}^{z(2)}\right), \\ \mathcal{B}^{i''j''} &= \frac{1}{\sqrt{3}}\epsilon^{i''j''}\left(\frac{2\sqrt{3}}{3}\mathcal{B}^z - 2\mathcal{B}^{z(2)}\right), \\ \mathcal{B}^{iz} &= -2\epsilon_j^i \mathcal{B}^j, \\ \mathcal{B}^{i'z} &= -2\epsilon_{j'}^{i'} \mathcal{B}^{j'}, \\ \mathcal{B}^{i''z} &= -2\epsilon_{j''}^{i''} \mathcal{B}^{j''}. \end{cases} \quad (2.11)$$

3 Harmonic analysis on Q^{111}

In this section we summarize the essential ideas concerning the techniques of harmonic analysis on homogeneous seven-manifolds originally developed in [19, 26]. These techniques are the basic ingredient of our calculations and in the present summary we present them already applied to the specific case of the Q^{111} manifold.

The essential goal of harmonic analysis is that of translating a differential equation problem into a linear algebraic one, by means of group theory. In the present case, the differential equations to solve are the linearized field equations of Kaluza Klein supergravity, whose typical form is:

$$\left(\square_x^{[J_1 J_2]} + \boxtimes_y^{[\lambda_1 \lambda_2 \lambda_3]}\right) \Phi_{[\lambda_1 \lambda_2 \lambda_3]}^{[J_1 J_2]}(x, y) = 0, \quad (3.1)$$

where $\Phi_{[\lambda_1 \lambda_2 \lambda_3]}^{[J_1 J_2]}(x, y)$ is a field transforming in the irreducible representations $[J_1 J_2]$ of $SO(3, 2)$ and $[\lambda_1 \lambda_2 \lambda_3]$ of $SO(7)$, and depends both on the coordinates x of Anti-de Sitter space and on the coordinates y of G/H . $\square_x^{[J_1 J_2]}$ is the kinetic operator for a field of spin $[J_1 J_2]$ in four dimension while $\boxtimes_y^{[\lambda_1 \lambda_2 \lambda_3]}$ is the kinetic operator for a field of spin $[\lambda_1 \lambda_2 \lambda_3]$ in seven dimensions.

Now, the harmonics constitute a complete set of functions for the expansion of any $SO(7)$ -irreducible field over G/H , $\mathcal{Y}_{[\lambda_1 \lambda_2 \lambda_3]}^i(y)$. But their most important property is that they transform irreducibly under G , the group of isometries of the coset space. This group acts on $\mathcal{Y}_{[\lambda_1 \lambda_2 \lambda_3]}^i(y)$ through the so-called covariant Lie derivative (see eq.(2.25) of [19]):

$$\delta_\Lambda \mathcal{Y}_{[\lambda_1 \lambda_2 \lambda_3]}^i(y) = \mathcal{L}_\Lambda \mathcal{Y}_{[\lambda_1 \lambda_2 \lambda_3]}^i(y), \quad (3.2)$$

which satisfy the Lie algebra of the group G :

$$[\mathcal{L}_\Lambda, \mathcal{L}_\Sigma] = \mathcal{C}_{\Lambda\Sigma}^\Pi \mathcal{L}_\Pi. \quad (3.3)$$

Moreover, the operators $\mathcal{L}_\Lambda$ commute with the $SO(7)$ covariant derivative:

$$\mathcal{L}_\Lambda D \mathcal{Y}^i = D \mathcal{L}_\Lambda \mathcal{Y}^i, \quad (3.4)$$

where D is defined by

$$D = d + \mathcal{B}^{\alpha\beta} t_{\alpha\beta}, \quad (3.5)$$

and $(t_{\alpha\beta})^i_j$ are the generators of the $SO(7)$ irreducible representation $[\lambda_1 \lambda_2 \lambda_3]$ of $\mathcal{Y}^i$ (e.g. $(t_{\alpha\beta})^{\gamma\delta} = -\delta_{\alpha\beta}^{\gamma\delta}$ for the vector representation).

An important thing to note is that $H = U(1)' \times U(1)'' \times U(1)'''$ is necessarily a subgroup of $SO(7)$, whose natural embedding is given in terms of the following embedding of the algebra $\mathbb{H}$ into the adjoint representation of $SO(7)$:

$$(T_H)^\alpha_\beta = \mathcal{C}_H^\alpha{}_\beta,$$

$$(T_{Z^{(1)}})^\alpha_\beta = \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \epsilon^{ij} & 0 & 0 \\ 0 & 0 & -\epsilon^{i'j'} & 0 \\ 0 & 0 & 0 & 0 \end{pmatrix}, \quad (3.6)$$

$$(T_{Z^{(2)}})^\alpha_\beta = \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & e^{ij} & 0 & 0 \\ 0 & 0 & \epsilon^{i'j'} & 0 \\ 0 & 0 & 0 & -2\epsilon^{i''j''} \end{pmatrix}, \quad (3.7)$$

$$(T_{Z^{(3)}})^\alpha_\beta = \frac{1}{\sqrt{3}} \begin{pmatrix} 0 & 0 & 0 & 0 \\ 0 & \epsilon^{ij} & 0 & 0 \\ 0 & 0 & \epsilon^{i'j'} & 0 \\ 0 & 0 & 0 & \epsilon^{i''j''} \end{pmatrix}. \quad (3.8)$$

This means that the $SO(7)$ -indices of the various n -forms can be split in different subsets, each one transforming into an irreducible representation of H and the $SO(7)$ irreducible representations $[\lambda_1\lambda_2\lambda_3]$ break into the direct sum of H irreducible representations.

3.1 The zero-form

Every harmonic is defined by its H ($[Z^{(1)}, Z^{(2)}, Z^{(3)}]$) representation. The only representation into which the $[0, 0, 0]$ (i.e. the scalar) of $SO(7)$ breaks under H , is obviously the H -scalar representation.

Solving the constraints (2.2, 2.3, 2.4) we see that every harmonic is conveniently identified by the following labels:

$\mathbf{J}_3$	$\mathbf{J}'_3$	$\mathbf{J}''_3$
$k/2$	$k/2$	$k/2$

(3.9)

where J_3, J'_3, J''_3 denote the third components of the three isospin and $k \in \mathbf{Z}$ is an integer number.

The Kaluza Klein mass operator for the zero-form $\mathcal{Y}$ is given by

$$\boxtimes^{[000]}\mathcal{Y} \equiv D_\beta D^\beta \mathcal{Y} = D_\beta^H D^{H\beta} \mathcal{Y}. \quad (3.10)$$

The computation of its eigenvalues, on the G -representations, is immediate:

$$\boxtimes^{[000]}\mathcal{Y} \equiv M_{(0)^3} \mathcal{Y} = H_0 \mathcal{Y} \quad (3.11)$$

$$H_0 \equiv 32 \left[J(J+1) + J'(J'+1) + J''(J''+1) - \frac{k^2}{4} \right] \quad (3.12)$$

3.2 The one-form

The decomposition under H of the vector representation of $SO(7)$ is the following:

$$[1, 0, 0] \rightarrow [0, 0, 0] \oplus [i, i, i] \oplus [-i, -i, -i] \oplus [i, -i, -i] \oplus [-i, i, i] \oplus [0, 2i, -i] \oplus [0, -2i, i]. \quad (3.13)$$

where the three digits in the brackets $[p, q, r]$ denote the eigenvalues of the generators $Z^{(1)}$, $Z^{(2)}$ and $Z^{(3)}$. Every harmonic is then conveniently identified by the following labels:

$\mathbf{J}_3$	$\mathbf{J}'_3$	$\mathbf{J}''_3$
$k/2$	$k/2$	$k/2$
$k/2 - 1$	$k/2$	$k/2$
$k/2 + 1$	$k/2$	$k/2$
$k/2$	$k/2 - 1$	$k/2$
$k/2$	$k/2 + 1$	$k/2$
$k/2$	$k/2$	$k/2 - 1$
$k/2$	$k/2$	$k/2 + 1$

(3.14)

The Laplace Beltrami operator for the transverse one-form field $\mathcal{Y}^\alpha$ is given by

$$\boxtimes^{[100]} \mathcal{Y}^\alpha \equiv M_{(1)(0)^2} \mathcal{Y}^\alpha = 2D_\beta D^{[\beta} \mathcal{Y}^{\alpha]} = (D^\beta D_\beta + 24) \mathcal{Y}^\alpha, \quad (3.15)$$

where transversality of $\mathcal{Y}^\alpha$ means that $D_\alpha \mathcal{Y}^\alpha = 0$. From the decomposition $D_\alpha = D_\alpha^H + \mathbb{M}_\alpha$ we obtain:

$$\boxtimes^{[100]} \mathcal{Y}^\alpha = (D^{H\beta} D_\beta^H + 24) \mathcal{Y}^\alpha + \eta^{\gamma\delta} \left(2(\mathbb{M}_\gamma)^\alpha{}_\beta D_\delta^H + (\mathbb{M}_\gamma)^\alpha{}_\epsilon (\mathbb{M}_\delta)^\epsilon{}_\beta \right) \mathcal{Y}^\beta. \quad (3.16)$$

The matrix of this operator on the AdS_4 fields is given by

Column one to four:

$$\left| \begin{array}{cccc} H_0 + 48 & 8 * \sqrt{2}(a + k/2 + 1) & -8 * \sqrt{2}(a - k/2 + 1) & 8 * \sqrt{2}(b + k/2 + 1) \\ 16\sqrt{2}(a - k/2) & H_0 - 16k & 0 & 0 \\ -16\sqrt{2}(a + k/2) & 0 & H_0 + 16k & 0 \\ 16\sqrt{2}(b - k/2) & 0 & 0 & H_0 - 16k \\ -16\sqrt{2}(b + k/2) & 0 & 0 & 0 \\ 16\sqrt{2}(c - k/2) & 0 & 0 & 0 \\ -16\sqrt{2}(c + k/2) & 0 & 0 & 0 \end{array} \right| \quad (3.17)$$

Column five to seven:

$$\left| \begin{array}{ccc} -8 * \sqrt{2}(b - k/2 + 1) & 8 * \sqrt{2}(c + k/2 + 1) & -8 * \sqrt{2}(c - k/2 + 1) \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \\ H_0 + 16k & 0 & 0 \\ 0 & H_0 - 16k & 0 \\ 0 & 0 & H_0 + 16k \end{array} \right| \quad (3.18)$$

This matrix has the following eigenvalues:

$$\begin{aligned}
\lambda_1 &= H_0 \\
\lambda_2 = \lambda_3 &= H_0 - 16k \\
\lambda_4 = \lambda_5 &= H_0 + 16k \\
\lambda_6 &= H_0 + 24 + 4\sqrt{H_0 + 36} \\
\lambda_7 &= H_0 + 24 - 4\sqrt{H_0 + 36}
\end{aligned} \tag{3.19}$$

in terms of H_0 defined in (3.12)

Our actual operator (3.17, 3.18) contains the eigenvalues of $M_{(0)^3}$, which are *longitudinal* (hence *non-physical*) for the one-form.

The eigenvalue λ_1 in (3.19) is the longitudinal one, equal to the zero-form eigenvalue H_0 .

3.3 The two-form mass operator

Under the action of $H = U(1)' \times U(1)'' \times U(1)'''$ the 21 components of the $SO(7)$ two-form transform into the completely reducible representation:

$$\begin{aligned}
[1, 1, 0] \rightarrow & [0, 2i, 2i] \oplus [0, -2i, -2i] \oplus [i, -i, 2i] \oplus [-i, i, -2i] \oplus [-i, -i, 2i] \oplus \\
& \oplus [i, i, -2i] \oplus [2i, 0, 0] \oplus [-2i, 0, 0] \oplus [i, 3i, 0] \\
& \oplus [-i, -3i, 0] \oplus [-i, 3i, 0] \oplus [i, -3i, 0] \oplus 3[0, 0, 0], \\
& \oplus [i, i, i] \oplus [-i, -i, -i] \oplus [i, -i, -i] \oplus [-i, i, i] \oplus [0, 2i, -i] \oplus [0, -2i, i]. \tag{3.20}
\end{aligned}$$

Every harmonic is then conveniently identified by the following labels:

$\mathbf{J}_3$	$\mathbf{J}'_3$	$\mathbf{J}''_3$
$k/2 + 1$	$k/2 + 1$	$k/2$
$k/2 - 1$	$k/2 - 1$	$k/2$
$k/2 + 1$	$k/2$	$k/2 + 1$
$k/2 - 1$	$k/2$	$k/2 - 1$
$k/2$	$k/2 + 1$	$k/2 + 1$
$k/2$	$k/2 - 1$	$k/2 - 1$
$k/2 + 1$	$k/2 - 1$	$k/2$
$k/2 - 1$	$k/2 + 1$	$k/2$
$k/2 + 1$	$k/2$	$k/2 - 1$
$k/2 - 1$	$k/2$	$k/2 + 1$
$k/2$	$k/2 + 1$	$k/2 - 1$
$k/2$	$k/2 - 1$	$k/2 + 1$
$k/2$	$k/2$	$k/2$
$k/2$	$k/2$	$k/2$
$k/2$	$k/2$	$k/2$
$k/2 + 1$	$k/2$	$k/2$
$k/2 - 1$	$k/2$	$k/2$
$k/2$	$k/2 + 1$	$k/2$
$k/2$	$k/2 - 1$	$k/2$
$k/2$	$k/2$	$k/2 + 1$
$k/2$	$k/2$	$k/2 - 1$

(3.21)

The Laplace Beltrami operator for the transverse two-form field $\mathcal{Y}^{\alpha\beta}$, is given by

$$\boxtimes^{[110]} \mathcal{Y}^{[\alpha\beta]} \equiv M_{(1)^2(0)} \mathcal{Y}^{[\alpha\beta]} = 3D_\gamma D^{[\gamma} \mathcal{Y}^{\alpha\beta]} = (D^\gamma D_\gamma + 48) \mathcal{Y}^{[\alpha\beta]} - 4\mathcal{R}^{[\alpha}{}_{[\gamma}{}^{\beta]}{}_{\delta]} \mathcal{Y}^{[\gamma\delta]} \quad (3.22)$$

From the decomposition $D_\alpha \mathcal{Y}^{\beta\gamma} = D_\alpha^H \mathcal{Y}^{\beta\gamma} + (\mathbb{M}_\alpha)^\beta{}_\delta \mathcal{Y}^{\delta\gamma} + (\mathbb{M}_\alpha)^\gamma{}_\delta \mathcal{Y}^{\beta\delta}$ we obtain:

$$\begin{aligned} \boxtimes^{[110]} \mathcal{Y}^{[\alpha\beta]} = & \left\{ 48 \delta^{[\alpha}{}_{[\gamma}{}^{\beta]}{}_{\delta]} - 4\mathcal{R}^{[\alpha}{}_{[\gamma}{}^{\beta]}{}_{\delta]} + \right. \\ & \left. 2\eta^{\mu\nu} (\mathbb{M}_\mu)^{[\alpha}{}_{[\gamma} (\mathbb{M}_\nu)^{\beta]}{}_{\delta]} + 2\eta^{\mu\nu} (\mathbb{M}_\mu \mathbb{M}_\nu)^{[\alpha}{}_{[\gamma} \delta^{\beta]}{}_{\delta]} + 4\eta^{\mu\nu} (\mathbb{M}_\mu)^{[\alpha}{}_{[\gamma} \delta^{\beta]}{}_{\delta]} D_\nu^H \right\} \mathcal{Y}^{[\gamma\delta]}. \end{aligned} \quad (3.23)$$

This operators acts on AdS_4 fields as the following 21×21 matrix:
Column one to five:

$H_0 - 32k$	0	0	0	0
0	$H_0 + 32k$	0	0	0
0	0	$H_0 - 32k$	0	0
0	0	0	$H_0 + 32k$	0
0	0	0	0	$H_0 - 32k$
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
$2\sqrt{32}(b + \frac{k}{2} + 1)$	0	$2\sqrt{32}(c + \frac{k}{2} + 1)$	0	0
0	$-2\sqrt{32}(b - \frac{k}{2} + 1)$	0	$-2\sqrt{32}(c - \frac{k}{2} + 1)$	0
$-2\sqrt{32}(a + \frac{k}{2} + 1)$	0	0	0	$2\sqrt{32}(c + \frac{k}{2} + 1)$
0	$2\sqrt{32}(a - \frac{k}{2} + 1)$	0	0	0
0	0	$-2\sqrt{32}(a + \frac{k}{2} + 1)$	0	$-2\sqrt{32}(b + \frac{k}{2} + 1)$
0	0	0	$2\sqrt{32}(a - \frac{k}{2} + 1)$	0

Column six to ten:

0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
$H_0 + 32k$	0	0	0	0
0	H_0	0	0	0
0	0	H_0	0	0
0	0	0	H_0	0
0	0	0	0	H_0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	$-2\sqrt{32}(b - \frac{k}{2} + 1)$	0	$-2\sqrt{32}(c - \frac{k}{2} + 1)$	0
0	0	$2\sqrt{32}(b + \frac{k}{2} + 1)$	0	$2\sqrt{32}(c + \frac{k}{2} + 1)$
0	0	$2\sqrt{32}(a - \frac{k}{2} + 1)$	0	0
$-2\sqrt{32}(c - \frac{k}{2} + 1)$	$-2\sqrt{32}(a + \frac{k}{2} + 1)$	0	0	0
0	0	0	0	$2\sqrt{32}(a - \frac{k}{2} + 1)$
$2\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	$-2\sqrt{32}(a + \frac{k}{2} + 1)$	0

Column eleven to sixteen:

0	0	0	0	0	$4\sqrt{32}(b - \frac{k}{2})$
0	0	0	0	0	0
0	0	0	0	0	$4\sqrt{32}(c - \frac{k}{2})$
0	0	0	0	0	0
0	0	0	0	0	0
0	0	0	0	0	0
0	0	0	0	0	$-4\sqrt{32}(b + \frac{k}{2})$
0	0	0	0	0	0
0	0	0	0	0	$-4\sqrt{32}(c + \frac{k}{2})$
0	0	0	0	0	0
H_0	0	0	0	0	0
0	H_0	0	0	0	0
0	0	$H_0 + 16$	16	16	$2i\sqrt{32}(a + \frac{k}{2} + 1)$
0	0	16	$H_0 + 16$	16	0
0	0	16	16	$H_0 + 16$	0
0	0	$-4i\sqrt{32}(a - \frac{k}{2})$	0	0	$H_0 + 32 - 16k$
0	0	$-4i\sqrt{32}(a + \frac{k}{2})$	0	0	0
$-2\sqrt{32}(c - \frac{k}{2} + 1)$	0	0	$-4i\sqrt{32}(b - \frac{k}{2})$	0	0
0	$2\sqrt{32}(c + \frac{k}{2} + 1)$	0	$-4i\sqrt{32}(b + \frac{k}{2})$	0	0
0	$2\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	$-4i\sqrt{32}(c - \frac{k}{2})$	0
$-2\sqrt{32}(b + \frac{k}{2} + 1)$	0	0	0	$-4i\sqrt{32}(c + \frac{k}{2})$	0

Column seventeen to twentyone:

0	$-4\sqrt{32}(a - \frac{k}{2})$	0	0	0
$-4\sqrt{32}(b + \frac{k}{2})$	0	$4\sqrt{32}(a + \frac{k}{2})$	0	0
0	0	0	$-4\sqrt{32}(a - \frac{k}{2})$	0
$-4\sqrt{32}(c + \frac{k}{2})$	0	0	0	$4\sqrt{32}(a + \frac{k}{2})$
0	$4\sqrt{32}(c - \frac{k}{2})$	0	$-4\sqrt{32}(b - \frac{k}{2})$	0
0	0	$-4\sqrt{32}(c + \frac{k}{2})$	0	$4\sqrt{32}(b + \frac{k}{2})$
0	0	$-4\sqrt{32}(a - \frac{k}{2})$	0	0
$4\sqrt{32}(b - \frac{k}{2})$	$4\sqrt{32}(a + \frac{k}{2})$	0	0	0
0	0	0	0	$-4\sqrt{32}(a - \frac{k}{2})$
$4\sqrt{32}(c - \frac{k}{2})$	0	0	$4\sqrt{32}(a + \frac{k}{2})$	0
0	$-4\sqrt{32}(c + \frac{k}{2})$	0	0	$-4\sqrt{32}(b - \frac{k}{2})$
0	0	$4\sqrt{32}(c - \frac{k}{2})$	$4\sqrt{32}(b + \frac{k}{2})$	0
$2i\sqrt{32}(a - \frac{k}{2} + 1)$	0	0	0	0
0	$2i\sqrt{32}(b + \frac{k}{2} + 1)$	$2i\sqrt{32}(b - \frac{k}{2} + 1)$	0	0
0	0	0	$2i\sqrt{32}(c + \frac{k}{2} + 1)$	$2i\sqrt{32}(c - \frac{k}{2} + 1)$
0	0	0	0	0
$H_0 + 32 + 16k$	0	0	0	0
0	$H_0 + 32 - 16k$	0	0	0
0	0	$H_0 + 32 + 16k$	0	0
0	0	0	$H_0 + 32 - 16k$	0
0	0	0	0	$H_0 + 32 + 16k$

This matrix has the following eigenvalues:

$$\lambda_1 = \lambda_2 \lambda_3 = H_0$$

$$\begin{aligned}
\lambda_4 = \lambda_5 &= H_0 - 16k \\
\lambda_6 = \lambda_7 &= H_0 + 16k \\
\lambda_8 &= H_0 + 24 + 4\sqrt{H_0 + 36} \\
\lambda_9 &= H_0 + 24 - 4\sqrt{H_0 + 36} \\
\lambda_{10} = \lambda_{11} &= H_0 + 32 \\
\lambda_{12} &= H_0 - 32k \\
\lambda_{13} &= H_0 + 32k \\
\lambda_{14} = \lambda_{15} &= H_0 + 16k + 16 + 4\sqrt{H_0 + 16k + 16} \\
\lambda_{16} = \lambda_{17} &= H_0 + 16k + 16 - 4\sqrt{H_0 + 16k + 16} \\
\lambda_{18} = \lambda_{19} &= H_0 - 16k + 16 - 4\sqrt{H_0 - 16k + 16} \\
\lambda_{20} = \lambda_{21} &= H_0 - 16k + 16 + 4\sqrt{H_0 - 16k + 16}
\end{aligned} \tag{3.24}$$

The eigenvalues $\lambda_1, \lambda_4, \lambda_5, \lambda_6, \lambda_7, \lambda_8, \lambda_9$ are those associated with longitudinal two forms that are non physical since they can be removed by gauge transformations. The remaining eigenvalues, associated with transverse two forms, provide the spectrum of physical states.

3.4 The three-form mass operator

The H decomposition of the three form into H irreducible fragments is the following:

$$\begin{aligned}
[1, 1, 1] \rightarrow & [0, 2i, 2i] \oplus 3[0, 0, 0] \oplus [0, 2i, 2i] \oplus [2i, 0, 0] \oplus [i, -i, 2i] \oplus \\
& \oplus [i, 3i, 0] \oplus [-2i, 0, 0] \oplus [0, -2i, -2i] \oplus [-i, -3i, 0] \oplus [-i, i, -2i] \oplus [-i, -i, 2i] \\
& \oplus [-i, 3i, 0] \oplus 3[i, -3i, 0] \oplus [i, i, -2i] \oplus [0, 0, 3i] \oplus [0, 4i, i] \\
& \oplus [2i, -2i, i] \oplus [2i, 2i, -i] \oplus [-2i, -2i, i] \oplus 2[i, i, i] \oplus 2[-i, -i, -i] \\
& \oplus 2[-i, i, i] \oplus 2[i, -i, -i] \oplus 2[0, -2i, i] \oplus 2[0, 2i, -i] \oplus [-2i, 2i, -i] \\
& \oplus [0, -4i, -i] \oplus [0, 0, -3i]
\end{aligned} \tag{3.25}$$

Every harmonic is then conveniently identified by the following labels:

$\mathbf{J}_3$	$\mathbf{J}'_3$	$\mathbf{J}''_3$
$k/2$	$k/2$	$k/2$
$k/2 + 1$	$k/2 + 1$	$k/2$
$k/2 + 1$	$k/2 - 1$	$k/2$
$k/2 + 1$	$k/2$	$k/2 + 1$
$k/2 + 1$	$k/2$	$k/2 - 1$
$k/2 - 1$	$k/2 + 1$	$k/2$
$k/2 - 1$	$k/2 - 1$	$k/2$
$k/2 - 1$	$k/2$	$k/2 + 1$
$k/2 - 1$	$k/2$	$k/2 - 1$
$k/2$	$k/2$	$k/2$
$k/2$	$k/2 + 1$	$k/2 + 1$
$k/2$	$k/2 + 1$	$k/2 - 1$
$k/2$	$k/2 - 1$	$k/2 + 1$
$k/2$	$k/2 - 1$	$k/2 - 1$
$k/2$	$k/2$	$k/2$
$k/2$	$k/2 + 1$	$k/2$
$k/2$	$k/2 - 1$	$k/2$
$k/2$	$k/2$	$k/2 + 1$
$k/2$	$k/2$	$k/2 - 1$
$k/2 + 1$	$k/2$	$k/2$
$k/2 + 1$	$k/2 + 1$	$k/2 + 1$
$k/2 + 1$	$k/2 + 1$	$k/2 - 1$
$k/2 + 1$	$k/2 - 1$	$k/2 + 1$
$k/2 + 1$	$k/2 - 1$	$k/2 - 1$
$k/2 + 1$	$k/2$	$k/2$
$k/2 - 1$	$k/2$	$k/2$
$k/2 - 1$	$k/2 + 1$	$k/2 + 1$
$k/2 - 1$	$k/2 + 1$	$k/2 - 1$
$k/2 - 1$	$k/2 - 1$	$k/2 + 1$
$k/2 - 1$	$k/2 - 1$	$k/2 - 1$
$k/2 - 1$	$k/2$	$k/2$
$k/2$	$k/2$	$k/2 + 1$
$k/2$	$k/2$	$k/2 - 1$
$k/2$	$k/2 + 1$	$k/2$
$k/2$	$k/2 - 1$	$k/2$

(3.26)

The Laplace Beltrami operator for the transverse three-form $\mathcal{Y}^{[\alpha\beta\gamma]}$, is a first-order differential operator, given by

$$\boxtimes^{[111]} \mathcal{Y}^{[\alpha\beta\gamma]} \equiv M_{(1)^3} \mathcal{Y}^{[\alpha\beta\gamma]} = \frac{1}{24} \epsilon^{\alpha\beta\gamma\delta}{}_{\mu\nu\rho} D_\delta \mathcal{Y}^{\mu\nu\rho} =$$

$$= \frac{1}{24} \epsilon^{\alpha\beta\gamma\delta}{}_{\mu\nu\rho} \left[D_\delta^H \mathcal{Y}^{\mu\nu\rho} + (\mathbb{M}_\delta)^\mu{}_\sigma \mathcal{Y}^{\sigma\nu\rho} + (\mathbb{M}_\delta)^\nu{}_\sigma \mathcal{Y}^{\mu\sigma\rho} + (\mathbb{M}_\delta)^\rho{}_\sigma \mathcal{Y}^{\mu\nu\sigma} \right]. \quad (3.27)$$

This operator acts on the AdS_4 fields as a 35×35 matrix:
Column one to six:

[illegible]

Column seven to eleven:

0	0	0	4	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
4	0	0	0	0	
0	-4	0	0	0	
0	0	4	0	0	
0	0	0	0	0	
0	0	0	0	4	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	4	0	
0	0	0	0	$-\frac{1}{2}\sqrt{32}(c + \frac{k}{2} + 1)$	
0	0	0	0	0	
0	0	0	$-i\sqrt{32}(c - \frac{k}{2})$	$\frac{1}{2}\sqrt{32}(b + \frac{k}{2} + 1)$	(3.29)
0	0	0	$i\sqrt{32}(c + \frac{k}{2})$	0	
0	0	0	0	0	
0	0	0	0	$-\sqrt{32}(a - \frac{k}{2})$	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	$-i\sqrt{32}(a - \frac{k}{2})$	0	
0	$\frac{1}{2}\sqrt{32}(c + \frac{k}{2} + 1)$	$-\frac{1}{2}\sqrt{32}(c - \frac{k}{2} + 1)$	0	0	
0	$-\sqrt{32}(b - \frac{k}{2})$	0	0	$\sqrt{32}(a + \frac{k}{2})$	
0	0	$\sqrt{32}(b - \frac{k}{2})$	0	0	
$-\sqrt{32}(c - \frac{k}{2})$	$\sqrt{32}(b + \frac{k}{2})$	0	0	0	
$\sqrt{32}(c + \frac{k}{2})$	0	$-\sqrt{32}(b + \frac{k}{2})$	0	0	
$-\frac{1}{2}\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	$i\sqrt{32}(a + \frac{k}{2})$	0	
0	$-\frac{1}{2}\sqrt{32}(a - \frac{k}{2} + 1)$	0	0	0	
0	0	$\frac{1}{2}\sqrt{32}(a - \frac{k}{2} + 1)$	0	0	
0	0	0	0	0	
$\frac{1}{2}\sqrt{32}(a - \frac{k}{2} + 1)$	0	0	0	0	

Column twelve to sixteen:

0	0	0	4	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	4	0
-4	0	0	0	$-\sqrt{32}(c - \frac{k}{2})$
0	-4	0	0	$\sqrt{32}(c + \frac{k}{2})$
0	0	4	0	0
0	0	0	0	$\frac{1}{2}i\sqrt{32}(b + \frac{k}{2} + 1)$
$\frac{1}{2}\sqrt{32}(c - \frac{k}{2} + 1)$	0	0	$-i\sqrt{32}(b - \frac{k}{2})$	0
0	$\frac{1}{2}\sqrt{32}(c + \frac{k}{2} + 1)$	$-\frac{1}{2}\sqrt{32}(c - \frac{k}{2} + 1)$	$i\sqrt{32}(b + \frac{k}{2})$	0
0	$-\frac{1}{2}\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	0
$-\frac{1}{2}\sqrt{32}(b + \frac{k}{2} + 1)$	0	$\frac{1}{2}\sqrt{32}(b - \frac{k}{2} + 1)$	0	0
0	0	0	$-i\sqrt{32}(a - \frac{k}{2})$	0
0	0	0	0	0
$\sqrt{32}(a - \frac{k}{2})$	0	0	0	0
0	$\sqrt{32}(a - \frac{k}{2})$	0	0	0
0	0	$-\sqrt{32}(a - \frac{k}{2})$	0	0
0	0	0	0	0
0	0	0	$i\sqrt{32}(a + \frac{k}{2})$	0
0	0	0	0	0
$-i\sqrt{32}(a + \frac{k}{2})$	0	0	0	0
0	$-\sqrt{32}(a + \frac{k}{2})$	0	0	0
0	0	$\sqrt{32}(a + \frac{k}{2})$	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	-4k
0	0	0	0	0

(3.30)

Column seventeen to twentyone:

0	0	0	0	0
0	0	0	0	$-\frac{1}{2}\sqrt{32}(c + \frac{k}{2} + 1)$
0	0	0	0	0
0	0	0	$-\sqrt{32}(c - \frac{k}{2})$	$\frac{1}{2}\sqrt{32}(b + \frac{k}{2} + 1)$
0	0	0	$\sqrt{32}(c + \frac{k}{2})$	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	$\frac{1}{2}i\sqrt{32}(c + \frac{k}{2} + 1)$	$-\frac{1}{2}i\sqrt{32}(c - \frac{k}{2} + 1)$	0	0
0	$\sqrt{32}(b - \frac{k}{2})$	0	0	$-\frac{1}{2}\sqrt{32}(a + \frac{k}{2} + 1)$
0	0	$-\sqrt{32}(b - \frac{k}{2})$	0	0
$\sqrt{32}(c - \frac{k}{2})$	$-\sqrt{32}(b + \frac{k}{2})$	0	0	0
$-\sqrt{32}(c + \frac{k}{2})$	0	0	0	0
$-\frac{1}{2}i\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	$\frac{1}{2}i\sqrt{32}(a + \frac{k}{2} + 1)$	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0
0	0	0	0	0

Column twentytwo to twentysix:

0	0	0	0	0	
$\frac{1}{2}\sqrt{32}(c - \frac{k}{2} + 1)$	0	0	$-\sqrt{32}(b - \frac{k}{2})$	0	
0	$\frac{1}{2}\sqrt{32}(c + \frac{k}{2} + 1)$	$-\frac{1}{2}\sqrt{32}(c - \frac{k}{2} + 1)$	$\sqrt{32}(b + \frac{k}{2})$	0	
0	$-\frac{1}{2}\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	0	
$-\frac{1}{2}\sqrt{32}(b + \frac{k}{2} + 1)$	0	$\frac{1}{2}\sqrt{32}(b - \frac{k}{2} + 1)$	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	$\sqrt{32}(c - \frac{k}{2})$	
0	0	0	0	$-\sqrt{32}(c + \frac{k}{2})$	
0	0	0	$\frac{1}{2}i\sqrt{32}(a + \frac{k}{2} + 1)$	0	
0	0	0	0	0	
$\frac{1}{2}\sqrt{32}(a + \frac{k}{2} + 1)$	0	0	0	0	
0	$\frac{1}{2}\sqrt{32}(a + \frac{k}{2} + 1)$	0	0	0	
0	0	$-\frac{1}{2}\sqrt{32}(a + \frac{k}{2} + 1)$	0	0	
0	0	0	0	$-\frac{1}{2}i\sqrt{32}(a - \frac{k}{2} + 1)$	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	
0	0	0	0	0	

Column twentyseven to thirtyone:

[illegible]

Column thirtytwo to thirtyfive:

[illegible]

This matrix has fifteen null eigenvalues corresponding to the longitudinal three-forms ($\mathcal{Y}^{(3)} = D \wedge \mathcal{Y}^{(2)}$), which are annihilated by $\boxtimes^{[111]}$ ($= *D \wedge$). These are gauge degrees of freedom that can be removed.

The remaining non-vanishing eigenvalues, associated with physical states, are:

$$\begin{aligned}\lambda_1 = \lambda_2 &= \frac{1}{4}\sqrt{H_0 + 16k + 16} \\ \lambda_3 = \lambda_4 &= -\frac{1}{4}\sqrt{H_0 + 16k + 16} \\ \lambda_5 = \lambda_6 &= \frac{1}{4}\sqrt{H_0 - 16k + 16} \\ \lambda_7 = \lambda_8 &= -\frac{1}{4}\sqrt{H_0 - 16k + 16}\end{aligned}$$

$$\begin{aligned}
\lambda_9 &= -\frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 36} \\
\lambda_{10} &= -\frac{1}{2} - \frac{1}{4}\sqrt{H_0 + 36} \\
\lambda_{11} = \lambda_{12} = \lambda_{13} &= \frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 4} \\
\lambda_{14} = \lambda_{15} = \lambda_{16} &= \frac{1}{2} - \frac{1}{4}\sqrt{H_0 + 4} \\
\lambda_{17} &= \frac{1}{2} + \frac{1}{4}\sqrt{H_0 - 32k + 4} \\
\lambda_{18} &= \frac{1}{2} - \frac{1}{4}\sqrt{H_0 - 32k + 4} \\
\lambda_{19} &= \frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 32k + 4} \\
\lambda_{20} &= \frac{1}{2} - \frac{1}{4}\sqrt{H_0 + 32k + 4}
\end{aligned} \tag{3.35}$$

3.5 The spinor

Under the action of H the 8 components of the $SO(7)$ spinor transform into the completely reducible representation:

$$\begin{aligned}
\left[\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right] \rightarrow & \left[-\frac{1}{2}i, 0, -2i\right] \oplus \left[-\frac{1}{2}i, -i, i\right] \oplus \left[-\frac{3}{2}i, 0, 0\right] \oplus \left[\frac{1}{2}i, -i, -i\right] \oplus \left[\frac{1}{2}i, i, -i\right] \oplus \\
& \left[\frac{1}{2}i, 0, 2i\right] \oplus \left[-\frac{1}{2}i, i, i\right] \oplus \left[\frac{3}{2}i, 0, 0\right]
\end{aligned} \tag{3.36}$$

Every harmonic is then conveniently identified by the following labels:

$\mathbf{J_3}$	$\mathbf{J'_3}$	$\mathbf{J''_3}$
$(k-1)/2$	$(k-1)/2$	$(k+1)/2$
$(k-1)/2$	$(k+1)/2$	$(k-1)/2$
$(k-1)/2$	$(k-1)/2$	$(k-1)/2$
$(k-1)/2$	$(k+1)/2$	$(k+1)/2$
$(k+1)/2$	$(k-1)/2$	$(k+1)/2$
$(k+1)/2$	$(k+1)/2$	$(k-1)/2$
$(k+1)/2$	$(k-1)/2$	$(k-1)/2$
$(k+1)/2$	$(k+1)/2$	$(k+1)/2$

(3.37)

This operator acts on the AdS_4 fields as a eight $\times$ eight matrix:

Column one to three:

$$\begin{vmatrix} 4\sqrt{2}(-\frac{7}{24}\sqrt{2}-\frac{1}{3}\sqrt{2}(-\frac{1}{2}+3\frac{k}{2})) & 0 & 4\sqrt{2}(-\frac{1}{2}-c+\frac{k}{2}) \\ 0 & 4\sqrt{2}(-\frac{7}{24}\sqrt{2}-\frac{1}{3}\sqrt{2}(-\frac{1}{2}+3\frac{k}{2})) & 4\sqrt{2}(\frac{1}{2}+b-\frac{k}{2}) \\ 4\sqrt{2}(-\frac{1}{2}-c-\frac{k}{2}) & 4\sqrt{2}(\frac{1}{2}+b+\frac{k}{2}) & 4\sqrt{2}(\frac{7}{8}\sqrt{2}+\frac{1}{3}\sqrt{2}(-\frac{3}{2}+3\frac{k}{2})) \\ 4\sqrt{2}(\frac{1}{2}+b-\frac{k}{2}) & 4\sqrt{2}(\frac{1}{2}+c-\frac{k}{2}) & 0 \\ 4\sqrt{2}(-\frac{1}{2}-a+\frac{k}{2}) & 0 & 0 \\ 0 & 4\sqrt{2}(-\frac{1}{2}-a+\frac{k}{2}) & 0 \\ 0 & 0 & 4\sqrt{2}(-\frac{1}{2}-a+\frac{k}{2}) \\ 0 & 0 & 0 \end{vmatrix} \quad (3.38)$$

Column four to six:

$$\begin{vmatrix} 4\sqrt{2}(\frac{1}{2}+b+\frac{k}{2}) & 4\sqrt{2}(-\frac{1}{2}-a-\frac{k}{2}) & 0 \\ 4\sqrt{2}(\frac{1}{2}+c+\frac{k}{2}) & 0 & 4\sqrt{2}(-\frac{1}{2}-a-\frac{k}{2}) \\ 0 & 0 & 0 \\ 4\sqrt{2}(-\frac{7}{24}\sqrt{2}+\frac{1}{3}\sqrt{2}(\frac{1}{2}+3\frac{k}{2})) & 0 & 0 \\ 0 & 4\sqrt{2}(-\frac{7}{24}\sqrt{2}+\frac{1}{3}\sqrt{2}(\frac{1}{2}+3\frac{k}{2})) & 0 \\ 0 & 0 & 4\sqrt{2}(-\frac{7}{24}\sqrt{2}+\frac{1}{3}\sqrt{2}(\frac{1}{2}+3\frac{k}{2})) \\ 0 & 4\sqrt{2}(\frac{1}{2}+c+\frac{k}{2}) & 4\sqrt{2}(-\frac{1}{2}-b-\frac{k}{2}) \\ 4\sqrt{2}(-\frac{1}{2}-a+\frac{k}{2}) & 4\sqrt{2}(-\frac{1}{2}-b+\frac{k}{2}) & 4\sqrt{2}(-\frac{1}{2}-c+\frac{k}{2}) \end{vmatrix} \quad (3.39)$$

Column seven and eight:

$$\begin{vmatrix} 0 & 0 \\ 0 & 0 \\ 4\sqrt{2}(-\frac{1}{2}-a-\frac{k}{2}) & 0 \\ 0 & 4\sqrt{2}(-\frac{1}{2}-a-\frac{k}{2}) \\ 4\sqrt{2}(\frac{1}{2}+c-\frac{k}{2}) & 4\sqrt{2}(-\frac{1}{2}-b-\frac{k}{2}) \\ 4\sqrt{2}(-\frac{1}{2}-b+\frac{k}{2}) & 4\sqrt{2}(-\frac{1}{2}-c-\frac{k}{2}) \\ 4\sqrt{2}(-\frac{7}{24}\sqrt{2}-\frac{1}{3}\sqrt{2}(-\frac{1}{2}+3\frac{k}{2})) & 0 \\ 0 & 4\sqrt{2}(\frac{7}{8}\sqrt{2}-\frac{1}{3}\sqrt{2}(\frac{3}{2}+3\frac{k}{2})) \end{vmatrix} \quad (3.40)$$

This matrix has the following eigenvalues:

$$\begin{aligned} \lambda_1 = \lambda_2 &= -8 + \sqrt{H_0 + 24} \\ \lambda_3 = \lambda_4 &= -8 - \sqrt{H_0 + 24} \\ \lambda_5 &= -6 + \sqrt{H_0 - 16k + 28} \\ \lambda_6 &= -6 + \sqrt{H_0 + 16k + 28} \\ \lambda_7 &= -6 - \sqrt{H_0 - 16k + 28} \\ \lambda_8 &= -6 - \sqrt{H_0 + 16k + 28} \end{aligned} \quad (3.41)$$

4 Organizing the spectrum of $Osp(2|4) \times SU(2) \times SU(2)' \times SU(2)''$ multiplets

In this section we present the $Osp(2|4)$ multiplets that appear compactifying $D = 11$ supergravity on $AdS_4 \times Q^{11}$. They are obtained by using the Kaluza-Klein mass spectrum derived in the previous sections together with the mass relations in [26, 18] and the multiplet structure of the $\mathcal{N} = 2$ multiplets. The procedure that we use is exactly the same as that employed in [5], to which we refer the interested reader. We report the structure of the $\mathcal{N} = 2$ supermultiplets in appendix A. The irreducible supermultiplets of $Osp(2|4)$ occur into irreducible representations of the bosonic group

$$G' = SU(2) \times SU(2) \times SU(2) ,$$

whose interpretation is the *flavor group* on the conformal field theory side of the correspondence and it is the *gauge group* on the supergravity side. In any case the crucial information one needs to extract from harmonic analysis is precisely the G' representation assignment of the supermultiplets and the actual value of E_0 and y_0 .

In the following pages, for each type of $\mathcal{N} = 2$ multiplet we list the G' representations through which it occurs in the spectrum. Furthermore for every multiplet, we give the energy and hypercharge values E_0 and y_0 of the Clifford vacuum. From tables 1, 2, 3, 4, 5, 6, 7, 8, 9 it is straightforward to get the energies and the hypercharges of all other fields in each multiplet.

4.1 Long multiplets

There are long multiplets for the following G' representations:

$$\left\{ J > \frac{k}{2}, J' > \frac{k}{2}, J'' > \frac{k}{2} \right\} \quad (4.1)$$

We have:

- one long graviton multiplet with

$$h : E_0 = \frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 36}, y_0 = k \quad (4.2)$$

- four long gravitino multiplets (two χ^+ and two χ^-) with

$$\chi^+ : E_0 = -\frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 16k + 16}, y_0 = k - 1 \quad (4.3)$$

$$\chi^+ : E_0 = -\frac{1}{2} + \frac{1}{4}\sqrt{H_0 - 16k + 16}, y_0 = k + 1 \quad (4.4)$$

$$\chi^- : E_0 = \frac{3}{2} + \frac{1}{4}\sqrt{H_0 + 16k + 16}, y_0 = k - 1 \quad (4.5)$$

$$\chi^- : E_0 = \frac{3}{2} + \frac{1}{4}\sqrt{H_0 - 16k + 16}, y_0 = k + 1 \quad (4.6)$$

- one W long vector multiplet with:

$$W : E_0 = \frac{5}{2} + \frac{1}{4}\sqrt{H_0 + 36}, y_0 = k \quad (4.7)$$

- one A long vector multiplet with:

$$A : E_0 = -\frac{3}{2} + \frac{1}{4}\sqrt{H_0 + 36}, \quad y_0 = k \quad (4.8)$$

- one Z long vector multiplet with

$$Z : E_0 = \frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 4}, \quad y_0 = k \quad (4.9)$$

- One Z long vector multiplet with

$$Z : E_0 = \frac{1}{2} + \frac{1}{4}\sqrt{H_0 + 32k + 4}, \quad y_0 = k \quad (4.10)$$

4.2 Short multiplets

The expansion of a generic field contains only the harmonics whose H - and G -quantum numbers are such that the G representation, decomposed under H , contains the H representation of the field. This fact poses some constraints on the G -quantum numbers.

Depending on which constraints are satisfied by a certain G representation, only part of the harmonics is present, and only their corresponding four-dimensional fields appear in the spectrum. Then, in the G representations in which such field disappear, there is *multiplet shortening*. In the modern perspective of Kaluza Klein theory, the exact spectrum of the short multiplets is crucial. Hence the importance of analyzing this disappearance of harmonics with care. We give here all the *multiplet shortening* found for the Q^{111} manifold. As already stressed in the introduction many of them confirm the gauge theory predictions made in [1]. We find also new type of shortening of the gravitino multiplet.

We have:

- One short graviton multiplet for the following G' representations:

$$\left\{ J = \frac{k}{2}, \quad J' = \frac{k}{2}, \quad J'' = \frac{k}{2} \right\} \quad (4.11)$$

with

$$E_0 = k + 2, \quad y_0 = k \quad (4.12)$$

- One short gravitino multiplet (χ^+) for the following G' representations:

$$\left\{ J = \frac{k}{2}, \quad J' = \frac{k}{2} + 1, \quad J'' = \frac{k}{2} + 1 \right\} \quad (4.13)$$

with

$$E_0 = k + \frac{5}{2}, \quad y_0 = k + 1 \quad (4.14)$$

All the other are obtained from (4.13) by permuting the role of the three $SU(2)$ groups.

- One short gravitino multiplet (χ^+) for the following G' representations:

$$\left\{ J = \frac{k}{2} - 1, \quad J' = \frac{k}{2}, \quad J'' = \frac{k}{2} \right\} \quad (4.15)$$

with

$$E_0 = k + \frac{1}{2}, \quad y_0 = k - 1 \quad (4.16)$$

All the other are obtained from (4.15) by permuting the role of the three $SU(2)$ groups.

- One short gravitino multiplet (χ^-) for the following G' representations:

$$\left\{ J = \frac{k}{2} - 1, \quad J' = \frac{k}{2}, \quad J'' = \frac{k}{2} \right\} \quad (4.17)$$

with

$$E_0 = k + \frac{1}{2}, \quad y_0 = k - 1 \quad (4.18)$$

All the other are obtained from (4.17) by permuting the role of the three $SU(2)$ groups.

- One short vector multiplet for the following G' representations:

$$\left\{ J = \frac{k}{2} + 1, \quad J' = \frac{k}{2}, \quad J'' = \frac{k}{2} \right\} \quad (4.19)$$

with

$$E_0 = k + 1, \quad y_0 = k \quad (4.20)$$

All the other are obtained from (4.19) by permuting the role of the three $SU(2)$ groups.

- One short vector multiplet for the following G' representations:

$$\left\{ J = \frac{k}{2}, \quad J' = \frac{k}{2}, \quad J'' = \frac{k}{2} \right\} \quad (4.21)$$

with

$$E_0 = k, \quad y_0 = k \quad (4.22)$$

- One hypermultiplet for the following G' representations:

$$\left\{ J = \frac{k}{2}, \quad J' = \frac{k}{2}, \quad J'' = \frac{k}{2} \right\} \quad (4.23)$$

with

$$E_0 = k, \quad y_0 = k \quad k \geq 1 \quad (4.24)$$

4.3 Massless multiplets

We have:

- the massless graviton multiplet in the singlet representation:

$$\{J = 0, J' = 0, J'' = 0\} \quad (4.25)$$

with

$$E_0 = 2, y_0 = 0 \quad (4.26)$$

- the massless vector multiplet in the adjoint representation:

$$\{J = 1, J' = 0, J'' = 0\} \quad (4.27)$$

with

$$E_0 = 1, y_0 = 0 \quad (4.28)$$

All the other are obtained from (4.27) by permuting the role of the three $SU(2)$ groups.

- two additional massless vector multiplets in the singlet representation:

$$\{J = 0, J' = 0, J'' = 0\} \quad (4.29)$$

with

$$E_0 = 1, y_0 = 0 \quad (4.30)$$

These two vector multiplets arises from the three form $A_{\mu ij}$ and are due to the existence of two closed cohomology two forms on the Q^{111} manifold. These multiplets are named the Betti multiplets.

5 Conclusions

In this paper I have verified that the SCFT conjectured in [1] to be the dual partner of M-theory compactified on Q^{111} is correct, since it reproduces the Kaluza-Klein spectrum of such a manifold. This result was expected on general grounds but a direct verification was so far missing. It was in my opinion important to make this final check since it supports the geometrical picture of the dual SCFT of $\mathcal{N} = 2 AdS_4 \times M^7$ compactifications based on Sasakian geometry.

Acknowledgement

I would like to thank D. Fabbri, L. Gualtieri, A. Zaffaroni and in particular P. Frè for very useful discussions.

A Structure of the $Osp(2|4)$ multiplets

The structure of a supermultiplet is conveniently described by listing the energy E , the spin s and the hypercharge y of all the fields of the multiplet.

In tables 1,2, 3,4, 5,6,7,8, 9 we provide such information in the first three columns. The remaining columns provide additional information concerning the way the abstract $Osp(2|4)$ supermultiplets are actually realized in Kaluza Klein supergravity. Indeed for each particle state appearing in the supermultiplet we write the name of the corresponding field in the Kaluza Klein expansion of $D = 11$ supergravity to which such a particle state contributes. For the standard expansion of linearized $D = 11$ supergravity and the conventions for the names of the $D = 4$ fields we refer the reader to [5].

The *long gravitino multiplet*, satisfying $E_0 > |y_0| + \frac{3}{2}$, has the structure displayed in table 2.

The *long vector multiplet*, satisfying $E_0 > |y_0| + 1$, has the structure displayed in table 3. Also the long vector multiplet has different realizations from the Kaluza Klein viewpoint.

The short multiplets are of two kinds: the *short graviton*, *gravitino* and *vector multiplets*, that saturate the bound

$$E_0 = |y_0| + s_0 + 1 \quad (\text{A.1})$$

and the *hypermultiplets* (spin 1/2 multiplets), that saturate the other bound

$$E_0 = |y_0| \quad \text{with} \quad |y_0| \geq \frac{1}{2}. \quad (\text{A.2})$$

The *short graviton multiplet*, satisfying $E_0 = |y_0| + 2$, has the structure displayed in table 4.

The *short gravitino multiplet*, satisfying $E_0 = |y_0| + \frac{3}{2}$, has the structure displayed in table 5.

The *short vector multiplet*, satisfying $E_0 = |y_0| + 1$, has the structure displayed in table 6.

We must stress that the multiplets displayed in tables 4, 5, 6 are only half of the story, since they can be viewed as the BPS states where $E_0 = y_0 + s_0 + 1$ and $y_0 > 0$. In addition one has also the anti BPS states. These are the short multiplets where $E_0 = -y_0 + s_0 + 1$ with $y_0 < 0$. The structure of these anti short multiplets can be easily read off from tables 4, 5, 6 by reversing the sign of all hypercharges.

The *hypermultiplet*, satisfying $E_0 = |y_0| \geq \frac{1}{2}$, has the structure displayed in table 7.

The massless multiplets are either short graviton or short vector multiplets satisfying the further condition

$$E_0 = s_0 + 1 \quad \text{equivalent to} \quad y_0 = 0. \quad (\text{A.3})$$

The *massless graviton multiplet*, satisfying $E_0 = 2$ $y_0 = 0$, has the structure displayed in table 8.

The *massless vector multiplet*, satisfying $E_0 = 1$ $y_0 = 0$, has the structure displayed in table 9. These are the $\mathcal{N} = 2$ supermultiplets in anti de Sitter space that can occur in Kaluza Klein supergravity.

References

- [1] Davide Fabbri et al., 3-D SUPERCONFORMAL THEORIES FROM SASAKIAN SEVEN MANIFOLDS: NEW NONTRIVIAL EVIDENCES FOR ADS(4) / CFT(3). Nucl. Phys. B577 (2000) 547-608. [hep-th/9907219](#)
- [2] Juan Maldacena, THE LARGE N LIMIT OF SUPERCONFORMAL FIELD THEORIES AND SUPERGRAVITY. Adv.Theor.Math.Phys.2:231-252,1998 [hep-th/9711200](#); for a review see: Ofer Aharony, Steven S. Gubser, Juan Maldacena, Hirosi Ooguri, Yaron Oz, LARGE N FIELD THEORIES, STRING THEORY AND GRAVITY. Phys. Rep. 323 (2000) 183-386. [hep-th/9905111](#)
- [3] S.S. Gubser, I.R. Klebanov, A.M. Polyakov, GAUGE THEORY CORRELATORS FROM NONCRITICAL STRING THEORY. Phys. Lett. B428 (1998) 105-114. [hep-th/9802109](#)
- [4] Edward Witten, ANTI-DE SITTER SPACE AND HOLOGRAPHY. Adv.Theor.Math.Phys.2:253-291,1998. [hep-th/9802150](#)
- [5] D. Fabbri, P. Fre', L. Gualtieri and P. Termonia, M-THEORY ON $AdS_4 \times M^{11}$: THE COMPLETE $Osp(2|4) \times SU(3) \times SU(2)$ SPECTRUM FROM HARMONIC ANALYSIS. [hep-th/9903036](#)
- [6] Peter G.O. Freund, Mark A. Rubin, DYNAMICS OF DIMENSIONAL REDUCTION. Phys.Lett.B97:233-235,1980.
- [7] M.J. Duff, B.E.W. Nilsson, C.N. Pope, KALUZA-KLEIN SUPERGRAVITY. Phys. Rep. 130 (1986) 1-142.
- [8] Kyungho Oh, Radu Tatar, THREE-DIMENSIONAL SCFT FROM M2- BRANES AT CONIFOLD SINGULARITIES. JHEP 9902:025,1999. [hep-th/9810244](#)
- [9] M.J. Duff, C.N. Pope, KALUZA-KLEIN SUPERGRAVITY AND THE SEVEN SPHERE. Trieste Workshop 1982:0183 (QC178:T7:1982) In *Trieste 1982, Proceedings, Supersymmetry and Supergravity '82*, 183-228 and London Imp. Coll. - ICTP-82- 83-07 (83,REC.MAR.) 49p.
- [10] M.A. Awada, M.J. Duff, C.N. Pope, N=8 SUPERGRAVITY BREAKS DOWN TO N=1. Phys.Rev.Lett.50:294,1983.
- [11] F. Englert, SPONTANEOUS COMPACTIFICATION OF ELEVEN- DIMENSIONAL SUPERGRAVITY. Phys.Lett.B119:339,1982.
- [12] B. Biran, F. Englert, B. de Wit, H. Nicolai, GAUGED N=8 SUPERGRAVITY AND ITS BREAKING FROM SPONTANEOUS COMPACTIFICATION. Phys.Lett.B124:45,1983., Erratum-ibid.B128:461,1983.
- [13] A. Casher, F. Englert, H. Nicolai, M. Roonan, THE MASS SPECTRUM OF SUPERGRAVITY ON THE ROUND SEVEN SPHERE. Nucl.Phys.B243:173,1984.

- [14] B. de Wit, H. Nicolai, ON THE RELATION BETWEEN $D = 4$ AND $D = 11$ SUPERGRAVITY. Nucl.Phys.B243:91,1984; B. de Wit, H. Nicolai, N.P. Warner, THE EMBEDDING OF GAUGED $N=8$ SUPERGRAVITY INTO $D = 11$ SUPERGRAVITY. Nucl.Phys.B255:29,1985; B. de Wit, H. Nicolai, THE CONSISTENCY OF THE S^{*7} TRUNCATION IN $D = 11$ SUPERGRAVITY. Nucl.Phys.B281:211,1987.
- [15] R. D'Auria, P. Fre, SPONTANEOUS GENERATION OF $OSP(4/8)$ SYMMETRY IN THE SPONTANEOUS COMPACTIFICATION OF $D = 11$ SUPERGRAVITY. Phys.Lett.B121:141,1983.
- [16] M. Gunaydin, N.P. Warner, UNITARY SUPERMULTIPLETS OF $OSP(8/4, R)$ AND THE SPECTRUM OF THE $S(7)$ COMPACTIFICATION OF ELEVEN-DIMENSIONAL SUPERGRAVITY. Nucl.Phys.B272:99,1986.
- [17] Edward Witten, SEARCH FOR A REALISTIC KALUZA-KLEIN THEORY. Nucl.Phys.B186:412,1981.
- [18] L. Castellani, R. D'Auria, P. Fre, $SU(3) \times SU(2) \times U(1)$ FROM $D = 11$ SUPERGRAVITY. Nucl.Phys.B239:610,1984.
- [19] R. D'Auria, P. Fre, ON THE FERMION MASS SPECTRUM OF KALUZA-KLEIN SUPERGRAVITY. Ann.Phys.157:1,1984.
- [20] R. D'Auria, P. Fre, ON THE SPECTRUM OF THE $N=2$ $SU(3) \times SU(2) \times U(1)$ GAUGE THEORY FROM $D = 11$ SUPERGRAVITY. Class.Quant.Grav.1:447,1984.
- [21] A. Ceresole, P. Fre, H. Nicolai, MULTIPLY STRUCTURE AND SPECTRA OF $N=2$ SUPERSYMMETRIC COMPACTIFICATIONS. Class.Quant.Grav.2:133,1985.
- [22] Leonardo Gualtieri, HARMONIC ANALYSIS AND SUPERCONFORMAL GAUGE THEORIES IN THREE-DIMENSIONS FROM ADS / CFT CORRESPONDENCE. ULB-TH-00-02 (Feb 1999) 219p. doctoral thesis. [hep-th/0002116](#)
- [23] Don N. Page, C.N. Pope, STABILITY ANALYSIS OF COMPACTIFICATIONS OF $D = 11$ SUPERGRAVITY WITH $SU(3) \times SU(2) \times U(1)$ SYMMETRY. Phys.Lett.B145:337,1984.
- [24] $N=2$ MATTER COUPLED SUPERGRAVITY FROM COMPACTIFICATION ON A COSET G/H POSSESSING AN ADDITIONAL KILLING VECTOR. Phys.Lett.B136:347,1984.
- [25] Don N. Page, C.N. Pope, WHICH COMPACTIFICATIONS OF $D = 11$ SUPERGRAVITY ARE STABLE? Phys.Lett.B144:346,1984.
- [26] R. D'Auria, P. Fre, UNIVERSAL BOSE-FERMI MASS RELATIONS IN KALUZA-KLEIN SUPERGRAVITY AND HARMONIC ANALYSIS ON COSET MANIFOLDS WITH KILLING SPINORS. Ann.Phys.162:372,1985.

- [27] L. Castellani, R. D'Auria, P. Fre, K. Pilch, P. van Nieuwenhuizen, THE BOSONIC MASS FORMULA FOR FREUND-RUBIN SOLUTIONS OF $D = 11$ SUPERGRAVITY ON GENERAL COSET MANIFOLDS. *Class.Quant.Grav.*1:339,1984.
- [28] Daniel Z. Freedman, Hermann Nicolai, MULTIPLY SHORTENING IN $OSP(N,4)$. *Nucl.Phys.B*237:342,1984.
- [29] L. Castellani, L.J. Romans, N.P. Warner, A CLASSIFICATION OF COMPACTIFYING SOLUTIONS FOR $D = 11$ SUPERGRAVITY. *Nucl.Phys.B*241:429,1984.
- [30] L. Castellani, R. D'Auria and P. Fre', SUPERGRAVITY AND SUPERSTRING THEORY: A GEOMETRIC PERSPECTIVE. World Scientific, Singapore, 1990.

Spin	Energy	Hypercharge	Mass (²)	Name
2	$E_0 + 1$	y_0	$16(E_0 + 1)(E_0 - 2)$	h
$\frac{3}{2}$	$E_0 + \frac{3}{2}$	$y_0 - 1$	$-4E_0 - 4$	χ^-
$\frac{3}{2}$	$E_0 + \frac{3}{2}$	$y_0 + 1$	$-4E_0 - 4$	χ^-
$\frac{3}{2}$	$E_0 + \frac{1}{2}$	$y_0 - 1$	$4E_0 - 8$	χ^+
$\frac{3}{2}$	$E_0 + \frac{1}{2}$	$y_0 + 1$	$4E_0 - 8$	χ^+
1	$E_0 + 2$	y_0	$16E_0(E_0 + 1)$	W
1	$E_0 + 1$	$y_0 - 2$	$16E_0(E_0 - 1)$	Z
1	$E_0 + 1$	$y_0 + 2$	$16E_0(E_0 - 1)$	Z
1	$E_0 + 1$	y_0	$16E_0(E_0 - 1)$	Z
1	$E_0 + 1$	y_0	$16E_0(E_0 - 1)$	Z
1	E_0	y_0	$16(E_0 - 1)(E_0 - 2)$	A
$\frac{1}{2}$	$E_0 + \frac{3}{2}$	$y_0 - 1$	$4E_0$	λ_T
$\frac{1}{2}$	$E_0 + \frac{3}{2}$	$y_0 + 1$	$4E_0$	λ_T
$\frac{1}{2}$	$E_0 + \frac{1}{2}$	$y_0 - 1$	$-4E_0 + 4$	λ_T
$\frac{1}{2}$	$E_0 + \frac{1}{2}$	$y_0 + 1$	$-4E_0 + 4$	λ_T
0	$E_0 + 1$	y_0	$16E_0(E_0 - 1)$	ϕ

Table 1: $\mathcal{N} = 2$ long graviton multiplet

Spin	Energy	Hypercharge	Mass (²)	Name	Mass (²)	name
$\frac{3}{2}$	$E_0 + 1$	y_0	$4E_0 - 6$	χ^+	$-4E_0 - 2$	χ^-
1	$E_0 + \frac{3}{2}$	$y_0 - 1$	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	Z	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	W
1	$E_0 + \frac{3}{2}$	$y_0 + 1$	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	Z	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	W
1	$E_0 + \frac{1}{2}$	$y_0 - 1$	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	A	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	Z
1	$E_0 + \frac{1}{2}$	$y_0 + 1$	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	A	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	Z
$\frac{1}{2}$	$E_0 + 2$	y_0	$4E_0 + 2$	λ_T	$-4E_0 - 2$	λ_L
$\frac{1}{2}$	$E_0 + 1$	$y_0 - 2$	$-4E_0 + 2$	λ_T	$-4E_0 - 2$	λ_T
$\frac{1}{2}$	$E_0 + 1$	y_0	$-4E_0 + 2$	λ_T	$4E_0 - 2$	λ_T
$\frac{1}{2}$	$E_0 + 1$	$y_0 + 2$	$-4E_0 + 2$	λ_T	$4E_0 - 2$	λ_T
$\frac{1}{2}$	$E_0 + 1$	y_0	$-4E_0 + 2$	λ_T	$4E_0 - 2$	λ_T
$\frac{1}{2}$	E_0	y_0	$4E_0 - 6$	λ_L	$-4E_0 + 6$	λ_T
0	$E_0 + \frac{3}{2}$	$y_0 - 1$	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	ϕ	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	π
0	$E_0 + \frac{3}{2}$	$y_0 + 1$	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	ϕ	$16(E_0 - \frac{1}{2})(E_0 + \frac{1}{2})$	π
0	$E_0 + \frac{1}{2}$	$y_0 - 1$	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	π	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	ϕ
0	$E_0 + \frac{1}{2}$	$y_0 + 1$	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	π	$16(E_0 - \frac{3}{2})(E_0 - \frac{1}{2})$	ϕ

Table 2: $\mathcal{N} = 2$ long gravitino multiplets χ^+ and χ^-

Spin	Energy	Hypercharge	Mass ⁽²⁾	Name	Name	Mass ⁽²⁾	Name
1	$E_0 + 1$	y_0	$16E_0(E_0 - 1)$	A	W	$16E_0(E_0 - 1)$	Z
$\frac{1}{2}$	$E_0 + \frac{3}{2}$	$y_0 - 1$	$-4E_0$	λ_T	λ_L	$4E_0$	λ_T
$\frac{1}{2}$	$E_0 + \frac{3}{2}$	$y_0 + 1$	$-4E_0$	λ_T	λ_L	$4E_0$	λ_T
$\frac{1}{2}$	$E_0 + \frac{1}{2}$	$y_0 - 1$	$4E_0 - 4$	λ_L	λ_T	$-4E_0 + 4$	λ_T
$\frac{1}{2}$	$E_0 + \frac{1}{2}$	$y_0 + 1$	$4E_0 - 4$	λ_L	λ_T	$-4E_0 + 4$	λ_T
0	$E_0 + 2$	y_0	$16E_0(E_0 + 1)$	ϕ	Σ	$16E_0(E_0 + 1)$	π
0	$E_0 + 1$	$y_0 - 2$	$16E_0(E_0 - 1)$	π	π	$16E_0(E_0 - 1)$	ϕ
0	$E_0 + 1$	$y_0 + 2$	$16E_0(E_0 - 1)$	π	π	$16E_0(E_0 - 1)$	ϕ
0	$E_0 + 1$	y_0	$16E_0(E_0 - 1)$	π	π	$16E_0(E_0 - 1)$	ϕ
0	E_0	y_0	$16(E_0 - 2)(E_0 - 1)$	S	ϕ	$16(E_0 - 2)(E_0 - 1)$	π

Table 3: $\mathcal{N} = 2$ long vector multiplets A, W and Z

Spin	Energy	Hypercharge	Mass ⁽²⁾	Name
2	$y_0 + 3$	y_0	$16y_0(y_0 + 3)$	h
$\frac{3}{2}$	$y_0 + \frac{7}{2}$	$y_0 - 1$	$-4y_0 - 12$	χ^-
$\frac{3}{2}$	$y_0 + \frac{7}{2}$	$y_0 + 1$	$4y_0$	χ^+
$\frac{3}{2}$	$y_0 + \frac{5}{2}$	$y_0 - 1$	$4y_0$	χ^+
1	$y_0 + 3$	$y_0 - 2$	$16(y_0 + 2)(y_0 + 1)$	Z
1	$y_0 + 3$	y_0	$16(y_0 + 2)(y_0 + 1)$	Z
1	$y_0 + 2$	y_0	$16y_0(y_0 + 1)$	A
$\frac{1}{2}$	$y_0 + \frac{5}{2}$	$y_0 - 1$	$-4y_0 - 4$	λ_T

Table 4: $\mathcal{N} = 2$ short graviton multiplet with positive hypercharge $y_0 > 0$

Spin	Energy	Hypercharge	Mass ⁽²⁾	Name
$\frac{3}{2}$	$y_0 + \frac{5}{2}$	y_0	$4y_0$	χ^+
1	$y_0 + 3$	$y_0 - 1$	$16(y_0 + 1)(y_0 + 2)$	Z
1	$y_0 + 2$	$y_0 + 1$	$16y_0(y_0 + 1)$	A
1	$y_0 + 2$	$y_0 - 1$	$16y_0(y_0 + 1)$	A
$\frac{1}{2}$	$y_0 + \frac{5}{2}$	y_0	$-4y_0 - 4$	λ_T
$\frac{1}{2}$	$y_0 + \frac{5}{2}$	$y_0 - 2$	$-4y_0 - 4$	λ_T
$\frac{1}{2}$	$y_0 + \frac{3}{2}$	y_0	$4y_0$	λ_L
0	$y_0 + 3$	$y_0 \pm 1$	$16(y_0 + 1)(y_0 + 2)$	ϕ

Table 5: $\mathcal{N} = 2$ short gravitino multiplet χ^+ with positive hypercharge $y_0 > 0$

Spin	Energy	Hypercharge	Mass ⁽²⁾	Name
1	$y_0 + 2$	y_0	$16y_0(y_0 + 1)$	A
$\frac{1}{2}$	$y_0 + \frac{5}{2}$	$y_0 \pm 1$	$-4y_0 - 4$	λ_T
$\frac{1}{2}$	$y_0 + \frac{3}{2}$	$y_0 + 1$	$4y_0$	λ_L
$\frac{1}{2}$	$y_0 + \frac{3}{2}$	$y_0 - 1$	$4y_0$	λ_L
0	$y_0 + 2$	$y_0 - 2$	$16y_0(y_0 + 1)$	π
0	$y_0 + 2$	y_0	$16y_0(y_0 + 1)$	π
0	$y_0 + 1$	y_0	$16y_0(y_0 - 1)$	S

Table 6: $\mathcal{N} = 2$ short vector multiplet A with positive hypercharge $y_0 > 0$

Spin	Energy	Hypercharge	Mass (²)	Name
$\frac{1}{2}$	$y_0 + \frac{1}{2}$	$y_0 - 1$	$4y_0 - 4$	λ_L
0	$y_0 + 1$	$y_0 - 2$	$16y_0(y_0 - 1)$	π
0	y_0	y_0	$16(y_0 - 2)(y_0 - 1)$	S
$\frac{1}{2}$	$y_0 + \frac{1}{2}$	$-y_0 + 1$	$4y_0 - 4$	λ_L
0	$y_0 + 1$	$-y_0 + 2$	$16y_0(y_0 - 1)$	π
0	y_0	$-y_0$	$16(y_0 - 1)(y_0 - 2)$	S

Table 7: $\mathcal{N} = 2$ hypermultiplet, $y_0 > 0$

Spin	Energy	Hypercharge	Mass (²)	Name
2	3	0	0	h
$\frac{3}{2}$	$\frac{5}{2}$	-1	0	χ^+
$\frac{3}{2}$	$\frac{5}{2}$	+1	0	χ^+
1	2	0	0	A

Table 8: $\mathcal{N} = 2$ massless graviton multiplet

Spin	Energy	Hypercharge	Mass (²)	Name	Mass (²)	Name
1	2	0	0	A	0	Z
$\frac{1}{2}$	$\frac{3}{2}$	-1	0	λ_L	0	λ_T
$\frac{1}{2}$	$\frac{3}{2}$	+1	0	λ_L	0	λ_T
0	2	0	0	π	0	ϕ
0	1	0	0	S	0	π

Table 9: $\mathcal{N} = 2$ massless vector multiplets A and Z